


Golden shiner (*Notemigonus crysoleucas*)

Habitat:

- feeding - lakes and impoundments and quiet pools of low gradient streams
- clear shallow water
- heavy vegetation


- spawning - vegetation



Emerald shiner (*Notropis atherinoides*)

Habitat:


- feeding - open-large stream channels and lake
 - low to moderate gradient
 - range of turbidities and bottom types
 - midwater or surface preferred, substrate of little importance
 - avoids rooted vegetation
- spawning - sand or firm mud substrate or gravel shoals



Blackchin shiner (*Notropis heterodon*)

Habitat:


- feeding - lakes, impoundments, and quiet pools in streams and rivers
- clear water
- clean sand, gravel, or organic debris substrate
- dense beds of submerged aquatic vegetation
- cannot tolerate turbidity, silt, or loss of aquatic vegetation



Blacknose shiner (*Notropis heterolepis*)

Habitat:


- feeding - clear lakes, impoundments, and pools of small, clear, low-gradient streams
 - aquatic vegetation
 - clean sand, gravel, marl, muck, peat, or organic debris substrate
 - cannot tolerate much turbidity, much siltation, or loss of aquatic vegetation
- spawning - sandy substrate



Spottail shiner (*Notropis hudsonius*)

Habitat:

- feeding - large rivers, lakes, and impoundments
 - firm sand and gravel substrate
 - low current
 - sparse to moderate vegetation
 - avoids turbidity
-
- spawning - over sandy shoals or gravelly riffles
 - near the mouths of small streams




Rosyface shiner (*Notropis rubellus*)

Habitat:

- feeding
 - moderate sized streams
 - moderate to high gradient
 - gravel or sand substrate; intolerant of silt substrate
 - clear water; intolerant of turbidity


- spawning
 - on nests of horneyhead chub, chesnut lamprey, and redhorses
 - sandy-gravel, gravel or bedrock substrate
 - shallow high gradient water



Sand shiner (*Notropis stramineus*)

Habitat:

- feeding - sand and gravel substrate
 - shallow pools in medium size streams, lakes, and impoundments
 - clear water and low gradient
 - rooted aquatic vegetation preferred
 - tolerant of some inorganic pollutants provided substrate is not covered
- spawning - clean gravel or sand substrate




Mimic shiner (*Notropis volucellus*)

Habitat:

- feeding - pools and backwater of streams, moderately weedy lakes and impoundments
- quiet or still water
- clear shallow water


spawning - aquatic vegetation necessary



Northern redbelly dace (*Phoxinus eos*)

Habitat:


- feeding - slow current
 - in boggy lakes and streams
 - detritus or silt substrate
 - clear to slightly turbid water
-
- spawning - filamentous algae needed for egg deposition



Bluntnose minnow (*Pimephales notatus*)

Habitat:

- feeding - quiet pools and backwaters of medium to large streams, lakes, and impoundments
 - clear warm water
 - some aquatic vegetation
 - firm substrates
 - tolerates all gradients, turbidity, organic and inorganic pollutants
-
- spawning - eggs deposited on the underside of flat stones or objects
 - nests in sand or gravel substrate




Fathead minnow (*Pimephales promelas*)

Habitat:

- feeding - pools of small streams, lakes, and impoundments
- tolerant of turbidity, high temperatures, and low oxygen


- spawning - on underside of objects in water 2 to 3 feet deep
- prefer sand, marl, or gravel substrate



Blacknose dace (*Rhinichthys atratulus*)

Habitat:


- feeding
 - moderate to high gradient streams
 - sand and gravel substrate
 - clear cool water in pools with deep holes and undercut banks
 - does not tolerate turbidity and silt well
- spawning - riffles with gravel substrate and fast current
- winter refuge - larger waters



Longnose dace (*Rhinichthys cataractae*)

Habitat:

- feeding - lakes and streams
- high gradient
- gravel or boulder substrate



Flint River Assessment Appendix


Creek chub (*Semotilus atromaculatus*)

Habitat:

- feeding - streams, rivers, or shore waters of lakes and impoundments
 - can tolerate intermittent flows
 - tolerates moderate turbidity

- spawning - gravel nests
 - low current

- winter refuge - deeper pools and runs



Quillback (*Carpoides cyprinus*)

Habitat:

- feeding - clear to turbid water
 - Lake Michigan
 - sand, sandy gravel, sandy silt, or clay-silt substrate
 - medium- to low-gradient rivers and streams; also lakes and sloughs
-
- spawning - streams or overflow areas of bends of rivers or bays of lakes
 - scatter eggs over sand or mud substrate





Figure 28.

Longnose sucker (*Catostomus catostomus*)

Habitat:

- feeding - clear, cold rivers and lakes

- spawning - in streams or lake shallows
 - current
 - gravel substrate




White sucker (*Catostomus commersoni*)

Habitat:

- feeding - streams, rivers, lakes, and impoundments
- can inhabit highly turbid and polluted waters

- spawning - quiet gravelly shallow areas of streams




Lake chubsucker (*Erimyzon sucetta*)

Habitat:

- feeding
 - larger clear streams, rivers, lakes, and impoundments
 - cannot tolerate turbid water
 - low gradient
 - prefers dense vegetation over substrate of sand or silt mixed with organic debris

- spawning
 - small clear streams with moderate to high gradient
 - sand or gravel substrate; no clayey silt




Northern hog sucker (*Hypentelium nigricans*)

Habitat:

- feeding - gravel or rubble substrate
- riffles and adjacent pools of warm shallow streams
- clear water
- doesn't like turbidity or siltation
- avoids profuse amounts of aquatic vegetation

- spawning - riffles
- shallow gravel substrate
- high gradient

- winter refuge - deeper quieter pools



Golden redhorse (*Moxostoma erythrurum*)

Habitat:

- feeding - warm medium gradient streams and rivers
 - clear riffly streams
 - medium size streams and rivers
 - tolerates some turbidity and silt
- spawning - shallow gravelly riffles
- winter refuge - larger streams

