



**STATE OF MICHIGAN  
DEPARTMENT OF NATURAL RESOURCES**

---

Number 21

June 1998

# **Manistee River Assessment Appendix**

Thomas J. Rozich

**MICHIGAN DEPARTMENT OF NATURAL RESOURCES  
FISHERIES DIVISION**

**Fisheries Special Report 21  
June 1998**

**Manistee River Assessment  
Appendix**

**Thomas J. Rozich**

The Michigan Department of Natural Resources, (MDNR) provides equal opportunities for employment and for access to Michigan's natural resources. State and Federal laws prohibit discrimination on the basis of race, color, sex, national origin, religion, disability, age, marital status, height and weight. If you believe that you have been discriminated against in any program, activity or facility, please write the MDNR Equal Opportunity Office, P.O. Box 30028, Lansing, MI 48909, or the Michigan Department of Civil Rights, 1200 6th Avenue, Detroit, MI 48226, or the Office of Human Resources, U.S. Fish and Wildlife Service, Washington D.C. 20204.

For more information about this publication or the American Disabilities Act (ADA), contact, Michigan Department of Natural Resources, Fisheries Division, Box 30446, Lansing, MI 48909, or call 517-373-1280.



*Printed under authority of Michigan Department of Natural Resources  
Total number of copies printed 375 — Total cost \$1453.93 — Cost per copy \$3.88*

Appendix I  
Distribution Maps of Fish Species


This appendix contains maps of known past and present fish distributions within the Manistee River watershed. The distributions of fish species were compiled from records located at the University of Michigan, Museums Fisheries Library, Michigan Department of Natural Resources, Institute for Fisheries Research, and offices in Cadillac and Grayling. Scientific names and phylogenic order follow Robins et al. (1991). Species that are listed under Michigan's Endangered Species Act (Part 365, Endangered Species Protection, of the Natural Resource and Environmental Protection Act, Act 451 of the Public Acts of 1994), their status follows their scientific name. Categories are declining, rare, threatened, endangered, extinct, and locally extinct.

Habitat descriptions were compiled from *The Fishes of Ohio* (Trautman 1982), *Freshwater Fishes of Canada* (Scott and Crossman 1973), *Fishes of Wisconsin* (Becker 1983), *Fishes of Missouri* (Pflieger 1975), and *Fishes of the Great Lakes Region* (Hubbs and Lagler 1947).

**Chestnut lamprey (*Ichthyomyzon castaneus*)**

**Habitat:**


- feeding - stable substrate of sand and silt with light growth of *chara* or quiet backwaters of muck and silt with dense rooted vegetation
- moderate current
- clear moderate-size water
  
- spawning - moderate-size stream
- nest builder



**Northern brook lamprey (*Ichthyomyzon fossor*)**

**Habitat:**


- feeding - young: low gradient, substrate with bars and beds of mixed sand and organic debris
- moderately warm water
  
- spawning - clear, high gradient streams (<15 feet wide)
- riffles with sand or gravel substrate



**American brook lamprey (*Lampetra appendix*)**

**Habitat:**


- feeding - young: low gradient, substrate with bars and beds of mixed sand and organic debris
  - clear cool stream water, sensitive to turbidity
  
- spawning - clear, high gradient streams (>15 feet wide)
  - cold water
  - gravel substrate
  
- winter refuge - sand or silt substrate for ammocoetes



**Sea lamprey (*Petromyzon marinus*)**

**Habitat:**


- feeding - young: substrate with beds of sand mixed with organic debris
- cannot tolerate silt
- adults: clear cool water of Lake Michigan
  
- spawning - no dams
- riffles with sand and gravel substrates



**Lake sturgeon (*Acipenser fulvescens*)**

**Habitat:**

- feeding - shoal areas of large rivers, lakes, and impoundments
- gravel, sand, rock substrates
  
- spawning - in or before rapids, at the base of dams in rivers
- in 2-15 feet of water
- swift current
- rocky ledges or around rocky islands in Great Lakes






**Longnose gar (*Lepisosteus osseus*)**

**Habitat:**


- feeding - adults: in deeper water
  - young: in shallows
  - clear water, low-gradient streams, lakes, and impoundments
  - will feed in moderate current
  - aquatic vegetation preferred, but not necessary
  - open water fish
- spawning - warm shallow water of lakes or streams over vegetation



**Bowfin** (*Amia calva*)

**Habitat:**


- feeding - clear water
  - abundant rooted aquatic vegetation
  - low gradient streams, lakes, and impoundments
  - tolerate only small amount of silt
  
- spawning - need vegetated water, 1 to 2 feet deep
  - can spawn under logs, stumps, or bushes
  
- winter refuge - gravelly pockets among aquatic vegetation



**Alewife** (*Alosa pseudoharengus*)

**Habitat:**


- feeding - adults: deep water of Lake Michigan
- young: shallow water of Lake Michigan
- prefers warmer waters
  
- spawning - streams or shallow beaches of lake
- sand or gravelly substrate
  
- winter refuge - deep water



**Gizzard shad (*Dorosoma cepedianum*)**

**Habitat:**


- feeding - large streams with low gradient, impoundments, and Lake Huron
- tolerant of clear and turbid water
  
- spawning - shallow areas of ponds, lakes, and large rivers
- low gradient



**Central stoneroller (*Campostoma anomalum*)**

**Habitat:**


- feeding - moderate to high gradients
  - rocky riffles
  - somewhat tolerant of turbidity
  - riffles and adjacent pools of warm, clear, shallow streams
  - gravel or cobble substrate
- spawning - riffles



**Lake chub (*Couesius plumbeus*) - rare**

**Habitat:**


- feeding - large rivers and lakes
- over a variety of substrates
  
- spawning - tributary streams
- rock substrate



**Spotfin shiner** (*Cyprinella spiloptera*)

**Habitat:**


- feeding
  - clear water tolerant of turbidity and siltation
  - some current
  - shallow depths
  - medium sized streams, lakes, and impoundments
  - clear sand or gravel substrate
  
- spawning
  - swift current
  - crevice spawner or on underside of submerged logs and roots



**Common carp (*Cyprinus carpio*)**

**Habitat:**

- feeding - low gradient fertile streams, rivers, lakes, and impoundments
  - abundance of aquatic vegetation or organic matter
  - tolerant of all substrates and clear to turbid water
- spawning - weedy or grassy shallows






**Brassy minnow** (*Hybognathus hankinsoni*)

**Habitat:**


- feeding - cool acidic streams
- slow to moderate current
- sand or gravel substrate



**Common shiner** (*Luxilus cornutus*)

**Habitat:**


- feeding - small, clear, high-gradient streams and rivers, or shores of clear water lakes and impoundments
  - gravel substrate
  - can tolerate some submerged aquatic vegetation
  - not very tolerant of turbidity or silted waters
  
- spawning - gravel nests of other fish, especially those at the head of a riffle



**Pearl dace (*Margariscus margarita*)**

**Habitat:**


- feeding - cool, neutral to acidic streams and lakes
- clear to slightly turbid water
  
- spawning - males are territorial
- clear water, 18-24 inches deep
- sand or gravel substrate
- weak to moderate current



**Hornyhead chub (*Nocomis biguttatus*)**

**Habitat:**


- feeding - adults: near riffles
  - young: near vegetation
  - clear water, does not tolerate turbidity
  - gravel substrate
  - low gradient streams that are tributaries to large streams
- 
- spawning - large stones and pebbles present
  - often below a riffle in shallow water
  - gravel substrate



**River chub** (*Nocomis micropogon*)

**Habitat:**


- feeding - moderate to large streams
- moderate to high gradient
- gravel, boulder, or bedrock substrate
- little to no aquatic vegetation
- cannot tolerate turbidity or siltation



**Golden shiner** (*Notemigonus crysoleucas*)

**Habitat:**


- feeding - lakes and impoundments and quiet pools of low gradient streams
- clear shallow water
- heavy vegetation
  
- spawning - vegetation



**Pugnose shiner (*Notropis anogenus*) - rare**

**Habitat:**


- feeding - very clear water of lakes, impoundments, and low-gradient streams
- aquatic vegetation
- clean sand, marl, or organic debris substrate
- extremely intolerant of turbidity



**Emerald shiner (*Notropis atherinoides*)**

**Habitat:**

- feeding - open-large stream channels and lake
  - low to moderate gradient
  - range of turbidites and bottom types
  - midwater or surface preferred, substrate of little importance
  - avoids rooted vegetation
- spawning - sand or firm mud substrate or gravel shoals






**Blackchin shiner (*Notropis heterodon*)**

**Habitat:**

- feeding - lakes, impoundments, and quiet pools in streams and rivers
- clear water
- clean sand, gravel, or organic debris substrate
- dense beds of submerged aquatic vegetation
- cannot tolerate turbidity, silt, or loss of aquatic vegetation




**Blacknose shiner** (*Notropis heterolepis*)

**Habitat:**

- feeding - clear lakes, impoundments, and pools of small, clear, low-gradient streams
- aquatic vegetation
- clean sand, gravel, marl, muck, peat, or organic debris substrate
- cannot tolerate much turbidity, much siltation, or loss of aquatic vegetation


spawning - sandy substrate



**Spottail shiner** (*Notropis hudsonius*)

**Habitat:**


- feeding - large rivers, lakes, and impoundments
- firm sand and gravel substrate
- low current
- sparse to moderate vegetation
- avoids turbidity
  
- spawning - over sandy shoals or gravelly riffles
- near the mouths of small streams



**Rosyface shiner** (*Notropis rubellus*)

**Habitat:**


- feeding - moderate sized streams
- moderate to high gradient
- gravel or sand substrate; intolerant of silt substrate
- clear water; intolerant of turbidity
  
- spawning - on nests of horneyhead chub, chesnut lamprey, and redhorses
- sandy-gravel, gravel or bedrock substrate
- shallow high gradient water



**Sand shiner** (*Notropis stramineus*)

**Habitat:**

- feeding - sand and gravel substrate
  - shallow pools in medium size streams, lakes, and impoundments
  - clear water and low gradient
  - rooted aquatic vegetation preferred
  - tolerant of some inorganic pollutants provided substrate is not covered
- spawning - clean gravel or sand substrate




**Mimic shiner** (*Notropis volucellus*)

**Habitat:**

- feeding - pools and backwater of streams, moderately weedy lakes and impoundments
- quiet or still water
- clear shallow water


- spawning - aquatic vegetation necessary



**Northern redbelly dace (*Phoxinus eos*)**

**Habitat:**


- feeding - slow current
  - in boggy lakes and streams
  - detritus or silt substrate
  - clear to slightly turbid water
- spawning - filamentous algae needed for egg deposition



**Finescale dace (*Phoxinus neogaeus*)**

**Habitat:**

- feeding - cool bog lakes and streams
- neutral to slightly acidic waters
- various substrates






**Bluntnose minnow (*Pimephales notatus*)**

**Habitat:**

- feeding - quiet pools and backwaters of medium to large streams, lakes, and impoundments
- clear warm water
- some aquatic vegetation
- firm substrates
- tolerates all gradients, turbidity, organic and inorganic pollutants


- spawning - eggs deposited on the underside of flat stones or objects
- nests in sand or gravel substrate



**Fathead minnow (*Pimephales promelas*)**

**Habitat:**

- feeding - pools of small streams, lakes, and impoundments
- tolerant of turbidity, high temperatures, and low oxygen
  
- spawning - on underside of objects in water 2 to 3 feet deep
- prefer sand, marl, or gravel substrate




**Blacknose dace (*Rhinichthys atratulus*)**

**Habitat:**

- feeding - moderate to high gradient streams
- sand and gravel substrate
- clear cool water in pools with deep holes and undercut banks
- does not tolerate turbidity and silt well

spawning - riffles with gravel substrate and fast current


winter refuge - larger waters



**Longnose dace (*Rhinichthys cataractae*)**

**Habitat:**


- feeding - lakes and streams
- high gradient
- gravel or boulder substrate



**Creek chub (*Semotilus atromaculatus*)**

**Habitat:**

- feeding - streams, rivers, or shore waters of lakes and impoundments
  - can tolerate intermittent flows
  - tolerates moderate turbidity
  
- spawning - gravel nests
  - low current
  
- winter refuge - deeper pools and runs




**Quillback (*Carpoides cyprinus*)**

**Habitat:**

- feeding - clear to turbid water
- Lake Michigan
- sand, sandy gravel, sandy silt, or clay-silt substrate
- medium- to low-gradient rivers and streams; also lakes and sloughs


- spawning - streams or overflow areas of bends of rivers or bays of lakes
- scatter eggs over sand or mud substrate



**Longnose sucker (*Catostomus catostomus*)**

**Habitat:**


- feeding - clear, cold rivers and lakes
  
- spawning - in streams or lake shallows
  - current
  - gravel substrate



**White sucker (*Catostomus commersoni*)**

**Habitat:**

- feeding - streams, rivers, lakes, and impoundments
- can inhabit highly turbid and polluted waters
  
- spawning - quiet gravelly shallow areas of streams






**Northern hog sucker (*Hypentelium nigricans*)**

**Habitat:**

- feeding - gravel or rubble substrate
- riffles and adjacent pools of warm shallow streams
- clear water
- doesn't like turbidity or siltation
- avoids profuse amounts of aquatic vegetation

- spawning - riffles
- shallow gravel substrate
- high gradient


- winter refuge - deeper quieter pools



**Silver redhorse (*Moxostoma anisurum*)**

**Habitat:**

- feeding - streams, rivers, lakes, and impoundments
- low current
- pollution and turbidity intolerant
  
- spawning - swift current in rivers, do not spawn in tributaries
- males territorial
- gravel to rubble substrate




**Golden redhorse (*Moxostoma erythrurum*)**

**Habitat:**

- feeding - warm medium gradient streams and rivers
- clear riffly streams
- medium size streams and rivers
- tolerates some turbidity and silt

spawning - shallow gravelly riffles


winter refuge - larger streams



**Shorthead redhorse** (*Moxostoma macrolepidotum*)

**Habitat:**


- feeding - downstream sections of large rivers, lakes, and impoundments
  - rocky substrates
  - swift water near riffles
  - clear to slightly turbid water
  
- spawning - gravelly riffles in smaller feeder streams



**Greater redhorse (*Moxostoma valenciennesi*)**

**Habitat:**


- feeding - large clear streams
  - clean sand, gravel, or boulder substrate
  - intolerant of excessive turbidity and chemical pollutants
  
- spawning - moderately rapid current



**Black bullhead (*Ameiurus melas*)**

**Habitat:**


- feeding - turbid water
  - silt bottom
  - low gradient small to medium streams, pools, and headwaters of large rivers; also in lakes and impoundments
  - can tolerate very warm water and very low dissolved oxygen
- spawning - nest in moderate to heavy vegetation or woody debris and under overhanging banks



**Yellow bullhead** (*Ameiurus natalis*)

**Habitat:**

- feeding - clear flowing water
  - heavy vegetation
  - low gradient streams, lakes, and impoundments
  - tolerant of low oxygen
- spawning - nest under a stream bank or near stones or stumps




**Brown bullhead** (*Ameiurus nebulosus*)

**Habitat:**

- feeding - larger streams and rivers, lakes and impoundments
- clear cool water with little clayey silt
- moderate amounts of aquatic vegetation
- sand, gravel, or muck substrate
- not tolerant of turbid water
- tolerant of warm water and low oxygen

- spawning - nest in mud or sand substrate among rooted aquatic vegetation usually near a stump, tree, or rock

- winter refuge - in muddy bottoms






**Channel catfish (*Ictalurus punctatus*)**

**Habitat:**


- feeding - moderately-clear, deeper waters of rivers, lakes, and impoundments
  - sand, gravel, or rubble substrate
  - low to moderate gradient
  
- spawning - secluded semi-dark areas such as holes, under banks, log jams, or rocks



**Tadpole madtom (*Noturus gyrinus*)**

**Habitat:**


- feeding
  - vegetative cover in low-moderate current waters
  - muddy substrate with extensive vegetation
  - clear waters of streams, rivers, and lakes
  
- spawning
  - mostly in rivers, sometimes shallows of lakes
  - nests in dark cavities (ex: beneath boards, logs, crayfish burrows)



**Northern pike** (*Esox lucius*)

**Habitat:**


- feeding - cool to moderately warm streams, rivers, lakes, and impoundments
- vegetation in slow to moderate current
  
- spawning - submerged vegetation with slow current in shallow water



**Tiger muskellunge** (*Esox masquinongy* x *E. lucius*)

**Habitat:**

- feeding - intermediate between muskellunge and northern pike
- spawning - hybrid species; muskellunge x northern pike
  - occasionally produced in wild, but most often from hatcheries
  - males are sterile, females may be fertile




**Muskellunge** (*Esox masquinongy*)

**Habitat:**

- feeding - warm, heavily vegetated lakes, stumpy weedy bays, and slow heavily vegetated medium to large rivers
- shallow cool water
- tolerant of low oxygen


spawning - clear shallow waters (15-20") in heavily vegetated areas



**Central mudminnow (*Umbra limi*)**

**Habitat:**


- feeding - undisturbed clear, low-gradient streams or rivers and lakes and impoundments
  - organic debris, muck, or peat substrates
  - aquatic vegetation
  
- spawning - floodplain areas, on vegetation



**Rainbow Smelt (*Osmerus mordax*)**

**Habitat:**

- feeding - midwater of lakes; 42-192 ft. in Lake Michigan
- spawning - in streams or off-shore shoals in Lake Michigan
  - gravel substrate
  - swift current




**Cisco {Lake herring} (*Coregonus artedii*)**

**Habitat:**

feeding - deep cool lakes, preferably oligotrophic

spawning - usually in lakes

- 3 to 6 feet of water with no vegetation
- often over gravel or stony substrate






**Lake Whitefish** (*Coregonus clupeaformis*)

**Habitat:**


- feeding - large cool water; Lake Michigan
- shallow water (for coregonids; 55-105 ft.)
  
- spawning - cold shallow water (<25 ft.)
- hard, stony, or sand substrate



**Pink salmon (*Oncorhynchus gorbuscha*)**

**Habitat:**


- feeding - large cold deep lakes - Lake Michigan
- spawning - gravel substrate in rivers
  - female prepares and guards nest until death



**Coho salmon** (*Oncorhynchus kisutch*)

**Habitat:**


- feeding - adults: Lake Michigan
- young: shallow gravel substrate in cold streams, later into pools
  
- spawning - cold streams and rivers
- swifter water of shallow gravelly substrate



**Rainbow trout (*Oncorhynchus mykiss*)**

**Habitat:**


- feeding - cold clear water of rivers and Lake Michigan
- moderate current
  
- spawning - gravelly riffles above a pool
- smaller tributaries



**Chinook salmon** (*Oncorhynchus tshawyscha*)

**Habitat:**

- feeding - adults: Lake Michigan
- young: shallow gravel substrate in cool streams, later into pools
  
- spawning - gravelly substrate in cool streams




**Round whitefish (*Prosopium cylindraceum*)**

**Habitat:**

feeding - lakes, rivers, and streams


spawning - shallows of lakes and rivers  
- gravel or rock substrate



**Brown trout (*Salmo trutta*)**

**Habitat:**


- feeding - cold, clear streams, rivers, and lakes (not >72°F)
  - medium to swift current in streams
  - does not tolerate silt well
  - prefers few individuals and species around
  - abundance of aquatic and land insects
- spawning - gravelly riffles; shallow headwater areas



**Brook trout (*Salvelinus fontinalis*)**

**Habitat:**

- feeding - cold, clear streams, rivers, and lakes (not >72°F)
- low current
- well oxygenated water
  
- spawning - gravelly riffles; shallow or headwater streams






**Splake** (*Salvelinus fontinalis* x *Salvelinus namaycush*)

**Habitat:**

- feeding - littoral habitat
- cool water lakes; also Lake Michigan
  
- spawning - hatchery produced cross of brook and lake trout
- offspring usually fertile, but with lower fecundity than either parent species




**Lake trout (*Salvelinus namaycush*)**

**Habitat:**

feeding - cold lakes and rivers


spawning - large boulder or rubble substrate  
- shallow water of lakes and rivers



**Trout-perch (*Percopsis omiscomaycus*)**

**Habitat:**


- feeding
  - clean sand or fine gravel substrate
  - long deep pools in low gradient streams and Lake Michigan
  - highly intolerant of clayey silts
  - avoids rooted aquatic vegetation
  
- spawning
  - over rocks in shallows
  - over sand and gravel substrates in Lake Michigan



**Burbot (*Lota lota*)**

**Habitat:**

- feeding - deep cold lakes and large cool rivers
  - mud, sand, rubble, boulder, silt, and gravel substrates
  
- spawning - in 1 to 4 feet of water in shallow bays or on shoals 5-10 feet deep usually in lakes, sometimes rivers
  - over sand or gravel substrate
  - under ice




**Banded killifish (*Fundulus diaphanus*)**

**Habitat:**

- feeding - quiet backwaters at the mouths of streams and lakes
- substrate of sand, gravel, and a few boulders
- also found over detritus substrate where patches of submerged aquatic vegetation are present


- spawning - quiet areas of weedy pools



**Brook silverside (*Labidesthes sicculus*)**

**Habitat:**


- feeding - clear, warm pools in streams and rivers; also lakes
  - does not tolerate turbidity
  - most frequently at surface
  
- spawning - in and around aquatic vegetation or over gravel substrate with a moderate current



**Brook stickleback (*Cluaea inconstans*)**

**Habitat:**


- feeding - clear, cold, densely vegetated streams, and swampy margins of lakes
  - low gradient
  - muck, peat, or marl substrate
  - not tolerant of turbidity
  
- spawning - shallow cool (<66°F) water
  - aquatic reeds or grasses necessary



**Ninespine stickleback (*Pungitius pungitius*)**

**Habitat:**

- feeding - open water of lakes; also Lake Michigan
- cool quiet waters
  
- spawning - builds nests among aquatic vegetation in creeks and streams






**Mottled sculpin (*Cottus bairdi*)**

**Habitat:**


- feeding - cool to cold streams
- riffle and rock substrates preferred
- clear to slightly turbid shallow water
  
- spawning - nests under logs or rock



**Slimy sculpin (*Cottus cognatus*)**

**Habitat:**


- feeding - cool lakes, impoundments, rivers, and streams
- gravel or rock substrate
  
- spawning - nest in shallow areas of lakes
- gravel substrate or rock ledge
- male parental care



**White bass (*Morone chrysops*)**

**Habitat:**


- feeding - large lakes, impoundments, and Lake Michigan
- clear water of 30 feet or less depth
- firm substrate
  
- spawning - tributary streams or shallow water of lakes
- over firm substrate



**Rock bass (*Ambloplites rupestris*)**

**Habitat:**


- feeding - clear, cool streams, rivers, and lakes
  - rocky to sand substrate
  - woody or vegetative cover
  
- spawning - sand or gravel nests
  - shallow water
  
- winter refuge - deep water



**Green sunfish (*Lepomis cyanellus*)**

**Habitat:**


- feeding - impoundments and lakes, and low-current streams and rivers
- no substrate preference
  
- spawning - nests in shallow areas sheltered by rocks, logs, or aquatic vegetation



**Pumpkinseed sunfish (*Lepomis gibbosus*)**

**Habitat:**


- feeding - non-flowing clear water in streams and rivers; also lakes and impoundments
  - muck or sand partly covered with organic debris substrate
  - dense beds of submerged aquatic vegetation
  
- spawning - nest in sand, gravel, or rock substrate
  - in shallow water near submerged vegetation



**Bluegill (*Lepomis macrochirus*)**

**Habitat:**


- feeding - non-flowing clear streams and rivers; also lakes and impoundments
  - sand, gravel, or muck containing organic debris substrate
  - scattered beds of aquatic vegetation
  - cannot tolerate low oxygen or continuous high turbidity and siltation
  
- spawning - nests in firm substrate of gravel, sand, or mud
  
- winter refuge - deep water



**Longear sunfish (*Lepomis megalotis*)**

**Habitat:**

- feeding - clear moderate-sized shallow streams with moderate vegetation
- rocky substrates
- little to no current
  
- spawning - nests in gravel, sand, or hard rock substrate






**Smallmouth bass (*Micropterus dolomieu*)**

**Habitat:**


- feeding - clear, cool, deep lakes and rivers
- streams where 40% consists of riffles over clean gravel, boulder, or bedrock substrate
- in pools with a current and >4 feet of depth
- gradients between 4 and 25 feet per mile
  
- spawning - nest in sandy, gravel, or rocky substrate
- gradients 7 to 25 feet per mile
- streams 20 to 100 feet wide
  
- winter refuge - larger deeper waters with gradients between 3 to 7 feet per mile



**Largemouth bass (*Micropterus salmoides*)**

**Habitat:**


- feeding - non-flowing clear waters - lakes, impoundments, and pools of streams
  - abundant aquatic vegetation
  - soft muck, organic debris, gravel, sand, and hard non-flocculent clay substrates
- 
- spawning - nest in gravelly sand to marl and soft mud substrates
  - emergent vegetation
  - quiet shallow bays; no current



**Black crappie (*Pomoxis nigromaculatus*)**

**Habitat:**


- feeding - larger clear non-silty low-gradient rivers; also in lakes and impoundments
  - clean hard sand or muck substrate
  - associated with submerged aquatic vegetation
  - does not tolerate silt or turbidity well
  
- spawning - nests in gravel, sand, or mud substrate
  - some vegetation must be present
  - sometimes nests under banks



**Rainbow darter (*Etheostoma caeruleum*)**

**Habitat:**


- feeding - gravelly high gradient riffles
- clear, moderate to large streams
- in shallows (average 1 foot)
  
- spawning - gravel or rubble riffles



**Iowa darter (*Etheostoma exile*)**

**Habitat:**


- feeding - clear, slow moving streams and lakes
  - sandy to muddy substrates
  - intolerant of turbid water
  - lives in rooted aquatic vegetation
  
- spawning - in pond-like extensions of streams on organic matter or roots
  - in shallows



**Johnny darter (*Etheostoma nigrum*)**

**Habitat:**


- feeding - sand and silt substrate
  - little to moderate current
  - shallow areas of streams, rivers, lakes, and impoundments
  - tolerant of many organic and inorganic pollutants and turbidity
- 
- spawning - underneath rocks
  - in stream pools or protected shallows of lakes



**Yellow perch (*Perca flavescens*)**

**Habitat:**


- feeding - clear lakes and impoundments; also Lake Michigan
- low gradient rivers
- abundance of rooted aquatics
- muck, organic debris, sand, or gravel substrate
- does not tolerate turbidity and siltation
  
- spawning - shallows of lakes, tributaries of streams
- occurs over rooted vegetation, submerged brush, fallen trees
- may occur over sand or gravel



**Logperch (*Percina caprodes*)**

**Habitat:**

- feeding - gravel riffles, deeper slower sections of rivers
  - medium size streams; also lakes, impoundments, and Lake Michigan
  - sand, gravel, or rock substrate
  - avoids turbidity and silt
- spawning - riffles or sandy in-shore shallows






**Blackside darter (*Percina maculata*)**

**Habitat:**


- feeding - small to medium streams
  - low to medium gradient
  - gravel and sand substrate
  - tolerate some turbidity
- spawning - gravel and sand substrate



**Walleye (*Stizostedion vitreum*)**

**Habitat:**

- feeding - larger, deeper streams and in large, shallow, turbid lakes and impoundments; also Lake Michigan
  - gravel, bedrock, and firm substrates preferred
  - does not tolerate a lot of turbidity or low oxygen
  
- spawning - rocky substrates in high gradient water in rivers
  - boulder to coarse gravel shoals in lakes
  
- winter refuge - avoids strong currents



**Freshwater drum (*Aplodinotus grunniens*)**

**Habitat:**

- feeding - deeper pools of rivers and Lake Michigan
- in shallows
- prefers clear waters and clean substrates
- can adapt to high turbidity levels

- spawning - pelagically, in open water, over sand or mud substrate
- occurs in bays or lower portions of marshes

