FSC-TPL-01-002 Application for a derogation to use a highly hazardous pesticide.

| 2,4-D, dma salt | |
|-----------------|--|
| | |

| Name and contact details of certification body requesting derogation: | SCS Dave Wager dwager@scscertified.com 510 251-7049 |
|---|---|
| Active ingredient for which derogation requested: | 2,4-D dma salt |
| Geographical scope of requested derogation: | Michigan |
| Is there an accredited or preliminarily accredited FSC Forest Stewardship Standard applicable to the territory concerned? | FSC US standard |
| Requested time period for derogation: (Derogations shall normally be issued for a five-year period. There will be a presumption against renewal of a derogation after the expiry of the five-year period). | 5 years |

Need may be demonstrated where:

- The pesticide is used for protecting native species and forests against damage caused by introduced species or for protecting human health against dangerous diseases, OR
- Use of the pesticide is obligatory under national laws or regulations, OR
 - Use of the pesticide is the only economically, environmentally, socially and technically feasible way of controlling specific organisms which are causing severe damage in natural forests or plantations in the specified country (as indicated by consideration, assessments and preferably field-trials of alternative nonchemical or less toxic pest-management methods)

Explain how the proposed use complies with the specified criteria for need, including the consideration of alternatives which do not require the use of pesticides on the FSC list of 'highly hazardous pesticides':

Overview

2,4-D dma salt is a selective herbicide used to control broad leaved plants such as woody species and forbs. 2,4-D dma salt is used to control non-native invasive plants in Michigan forests and openlands. Michigan openlands include grasslands, jack pine barrens, wildlife openings, roadways, and rights-of-way.

Specifics

More specifically, 2,4-D dma salt is used in combination with other herbicides or independently as part to the control of non-native invasives species such as honeylocust, honeysuckle, kudzu, multiflora rose, spotted knapweed, kochia, and Canada thistle. 2,4-D dma salt is a selective herbicide for broadleaf plants and does not control grasses. Using a product that is selective and affects as few plants as possible has been a preferred approach compared to unnecessarily applying a broader spectrum product.

In Michigan, invasive species such as those listed above reduce the biodiversity of our forest, reduce regeneration of important native trees, and reduce forest health. Michigan Department of Natural Resources (MDNR) manages over 4.5 million acres of diverse cover types that are mainly found in the northern Lower Peninsula and Upper Peninsula. The Draft 2006 State Forest Management Plan states that the introduction of non-native plant and animal species and diseases are a serious threat to the health of the State's forest ecosystems, and can have major ecological consequences for the composition of native forest communities. The desired future condition of state forest is that they be free from invasive plant and animal species that degrade ecological and socio-economic values and productivity, or the biological impact of such species is mitigated to the extent possible.

Michigan's Wildlife Action Plan (WAP) identifies threats to wildlife and landscape features that were evaluated as high severity throughout the State, one of these highest identified threats was the introduction of invasive non-native species.

As many as one-third of Michigan's plant species may now be non-native. In the Great Lakes basin, at least 37 terrestrial plant species and seven terrestrial insect species are invasive and pose threats to natural communities in Michigan.

Mechanical control or prescribed burring can sometimes be used to control invasive plants. However, these management practices may also increase rate of spread, germination or resprouting. Often these practices are used cooperatively or in conjunction with herbicide treatments.

There are numerous types of herbicides that can be used to control non-native invasive plants, including glyphosate. There are concerns that with repetitive treatments of specific herbicides such as glyphosate, that individual plants' will develop a tolerance to a specific chemical. By alternating chemical types and approaches there is less risk of the development of tolerance to specific chemicals.

2,4,-D dma salt is a selective herbicide that controls broadleaf plants and does not harm native grasses. Using a product that is selective and affects as few plants as possible has been a preferred approach compared to unnecessarily applying a broader spectrum product.

Not Controlling Invasive Plants

Invasive species left unchecked would compete with native plants, intercept sunlight, and monopolize available soil nutrients and moisture, resulting in slower growth of native plants, mortality and ultimately poor system health. Herbicide treatment of invasives is often a key step to control which, may also include mechanical treatments or prescribed fire. Furthermore, there are concerns that with repetitive treatments of specific treatments such as glyphosate those individual plants will develop a tolerance to a specific chemical. By alternating chemical types and approaches there is less risk of the development of tolerance to specific chemicals.

Control of annual, perennial and woody weeds is essential for the successful restoration, establishment and growth of native ecosystems. Without weed control, plants may die due to inability to compete for water and nutrients or growth rates may be so low that they can not compete against non-native plants. Effects can range from widespread mortality in new plantings to severe suppression of entire stands for indefinite periods.

2.4.-D & environment

2,4-D is one of the oldest herbicides used in the United States, first developed during World War II (TNC, 200X). Today, 2,4-D continues to be one of the most commonly used herbicides on the market. Because there is no longer a patent governing the manufacture and sale of 2,4-D, any company is free to produce it. Thus, a variety of inexpensive 2,4-D products are available from different manufacturers. Because it has been in use for so long, many of the studies regarding its behaviour in the environment are old (e.g. pre-1980). 2,4-D is a selective herbicide that kills dicots (but not grasses) by mimicking the growth hormone auxin, which causes uncontrolled growth and eventually death in susceptible plants. The half-life of 2,4-D in the environment is relatively short, averaging 10 days in soils and less than ten days in water, but can be significantly longer in cold, dry soils, or where the appropriate microbial community is not present to facilitate degradation. In the environment, most formulations are degraded to the anionic form, which is water-soluble and has the potential to be highly mobile. Ester formulations are toxic to fish and aquatic invertebrates, but salt formulations are registered for use against aquatic weeds. 2,4-D is of relatively low toxicity to animals but some formulations can cause severe eye damage. Certain crops, such as grapes, are highly sensitive to 2,4-D and application of this

herbicide should be avoided if they are nearby. Most formulations are highly volatile and should not be applied when conditions are windy or when temperatures are high.

The World Health Organization (1984) concluded that 2,4-D does not accumulate or persist in the environment. The primary degradation mechanism is microbial metabolism, but mineralization and possibly photolysis may also play a role. The average half-life (the time it takes for the herbicide concentration to decline by 50%) is 10 days, but rates of degradation can vary from several hours to several months or longer. Degradation rates are determined by the microbial population, environmental pH, soil moisture, and temperature (Que Hee & Sutherland 1981; Sandmann et al. 1988; Wilson et al. 1997). The type of 2,4-D formulation applied does not significantly affect the rate of degradation (Wilson et al. 1997).

2,4-D may be applied in acid, salt, or ester formulations, but in most cases, each of these formulations are apparently converted rapidly to the acid form once it contacts soil (Foster & McKercher 1973; Smith 1988; Wilson et al. 1997). Consequently, the rate of dissipation from soils is often the same regardless of the formulation of 2,4-D that is applied (Wilson et al. 1997). Half-lives are short, ranging from a few days to several months but detectable residues can persist for up to a year (McCall et al. 1981).

Degradation is almost entirely through microbial metabolism. Soil conditions that maximize microbial populations (i.e. warm and moist with a high organic content) maximize degradation rates (Foster & McKercher 1973; Ou 1984; Han & New 1994; Johnson et al. 1995a; Veeh et al. 1996).

Most formulations of 2,4-D do not bind tightly with soils and, therefore, have the potential to leach down into the soil column and to move off-site in surface or subsurface water flows. Leaching of 2,4-D to 30 cm has been reported (Johnson et al. 1995a). In many cases, extensive leaching does not occur, most likely because of the rapid degradation of the herbicide (Que Hee & Sutherland 1981). Where 2,4-D does leach, however, it will be more persistent because populations of microbes responsible for the degradation of 2,4-D tends to decrease with soil depth (Wilson et al. 1997).

2,4-D will change form and function with changes in water pH (Que Hee & Sutherland 1981). In alkaline (high pH; pH > 7) waters, 2,4-D takes an ionized (negatively charged) form that is water-soluble and remains in the water column. Theoretically, in water of a lower pH, 2,4-D will remain in a neutral molecular form, increasing its potential for adsorption to organic particles in water, and increasing its persistence (Wang et al. 1994a). 2,4-D is most likely to adsorb to suspended particles in muddy waters with a fine silt load (Que Hee & Sutherland 1981).

Que Hee and Sutherland (1981) reported that concentrations of most 2,4-D residues found in lakes and streams are < 1 ppm, although concentrations of up to 61 ppm have been reported immediately following direct application to water bodies. These concentrations are well above the 0.1 ppm established as "permissible" levels for potable water by the U.S. E.P.A. (EPA 1998).

2,4-D residues taken up by plants remain intact in the foliage until it is lost as litter and degraded in soils (Newton et al. 1990). Fruits from treated trees have been found to retain 2,4-D residues for up to seven weeks (Que Hee & Sutherland 1981).

- 2,4-D is considered of moderate toxicity to animals, although LD50 levels vary significantly between formulations and animal species (Ibrahim et al. 1991). The majority of LD50 values range between 300-1,000 mg/kg. For example, the LD50 for 2,4-D acid in rats and bobwhite quail is 764 mg/kg and 500 mg/kg, respectively. Some animals such as dogs, however, are significantly more sensitive to 2,4-D organic acids than are rats and humans (Ibrahim et al. 1991). In 1991, Hayes et al. reported a significant increase in the occurrence of malignant lymphoma among dogs whose owners applied 2,4-D to their lawns.
- 2,4-D can bio-accumulate in animals. In Russia, residues of more than ten times the allowable level were found in eggs, milk, and meat products served by public caterers and one study reported residues in 46% of tested cattle (Que Hee & Sutherland 1981). Risk to browsing wildlife, however, is low, Newton et al (1990) analysed 2,4-D residues in forest browse following aerial application to forests in Oregon and found them to be below the concentrations known to cause effects in mammals.

LC50 levels for bluegill sunfish and rainbow trout are 263 and 377 mg/L, respectively. Wang et al. (1994b) studied bioaccumulation of 2,4-D in carp and tilapia and found that accumulation of up to 18 times the ambient concentration occurred within two days of exposure. 2,4-D was found in oysters and clams in concentrations up to 3.8 ppm, and it persisted for up to two months (Thomas & Duffy 1968). The highest concentrations of 2,4-D were generally reached shortly after application, and dissipated within three weeks following exposure.

2,4-D can accumulate in fish exposed to concentrations as low as 0.05 ppm (Wang et al 1994b) and concentrations of 1.5 ppm can kill the eggs of fathead minnows in 48 hours (Thomas & Duffy 1968). After animals are removed from contaminated waters, they tend to excrete residues.

Human Safety

2,4-D can be absorbed through the skin or through the lungs if inhaled. Applicators of 2,4-D, particularly those using back-pack sprayers, are at greatest risk of exposure (Ibrahim et al. 1991; Johnson & Wattenberg 1996). Reported airborne residues of 1-35 micrograms/cubic meter of air when 2,4-D was applied using hand-held spray guns along power line right-of-ways. Absorption through the skin accounts for 90% of the 2,4-D absorbed by applicators (Ibrahim et al. 1991).

Once in the body, 2,4-D is distributed rapidly with the greatest concentrations appearing in the kidneys and liver (Johnson & Wattenberg 1996). The majority of the compound is excreted unmetabolized (Ibrahim et al. 1991). Due to its solubility in water, 2,4-D is not believed to accumulate in tissues, but is excreted in the urine in less than a week (Shearer 1980; Ibrahim et al. 1991; Johnson & Wattenberg 1996). Nevertheless, some agricultural workers and other applicators have experienced long term complications including pain, paresthesias (tingling or numbness), and paralysis following exposure to 2,4-D (Shearer 1980).

In 1991, a panel with expertise in epidemiology, toxicology, exposure assessment, and industrial hygiene convened to review the evidence available regarding the human carcinogenicity of 2,4-D (Ibrahim et al. 1991). The panel found that case-control studies showed evidence of a relationship between 2,4-D exposure and non-Hodgkins lymphoma in humans, with some studies showing an increased risk with increased exposure level (Ibrahim et al. 1991). Non-

Hodgkin's lymphoma is the human equivalent of the canine malignant lymphoma found to be associated with 2,4-D exposure in dogs (Hayes et al. 1991). When all evidence was evaluated, however, the panel could not find a cause-effect relationship between exposure to 2,4-D and human cancer (Ibrahim et al. 1991).

In another study of human exposure, female applicators were found to have a significant increase in cervical cancer associated with 2,4-D application. Due to the many confounding factors that make identification of cause and effect mechanisms difficult, other expert review panels including the U.S. EPA, Agriculture and Agri-food Canada, and the World Health Organization concluded that 2,4-D alone is not carcinogenic (Ibrahim et al. 1991; Mullison and Bond 1991).

2. Specified controls to mitigate the hazard

The derogation shall specify the controls that will be implemented to mitigate the hazard associated with the use of the pesticide, for example restrictions related to weather conditions, soil types, application method, water courses, etc..

If the specified formulation is considered to reduce the level of hazard then the information on which this claim is based shall be presented, and the applicant shall provide credible independent, third party support for the claimed reduction of hazard.

Specify the controls that will be implemented to mitigate the hazard:

Herbicides sold in the United States must be registered with the Federal government and in some cases by state regulatory agencies. They are reviewed and regulated by the U.S. Environmental Protection Agency (USEPA) under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA 1974; 7 J.S.C. 135 et seq., Public Laws 92-516, 94-140, and 95-356) and recent amendments. EPA regulations are enforced in Michigan through the Michigan Department of Agriculture.

The printed information and instructional material that is sold with a registered herbicide is known as the "label," and constitutes a legal document. These instructions are considered a part of compliance with FIFRA and other Federal regulations, and failure to use an herbicide in accord with label restrictions can lead to severe penalties. The label provides information on the chemical compound(s) comprising the active ingredient(s) of the herbicide, directions for correct use on target plant species, warnings and restrictions, and safety and antidote information.

Purchasers and applicators of restricted-use pesticides must comply with the certification requirements of the 1994 Michigan Natural Resources and Environmental Protection Act as amended (P.A. 451), Part 83 and detailed in Regulation 636 "Pesticide Applicators". This requires studying training manuals prepared by Michigan State University Extension and passing an examination administered by MDA. Recertification is required every three years and may be obtained by one of two methods. The applicator may study a training manual (Extension Bulletin E-2195) and pass an examination, or attend classes accredited by MDA for continuing education credits and obtain sufficient credits for the specific category of certification. Both methods ensure that additional information was provided to applicators in the safe and effective use of restricted-use pesticides.

Resource Application

As part of operational planning process alternatives are evaluated to control invasive plant species. Mechanical, biological, cultural as well as chemical treatments are evaluated for effectiveness and for cost efficiencies. In many cases several alternative control methods will need to be implemented for control of aggressive non-native invasive plants or populations. Frequently the application may be as a strip or spot application where as little as 10% to 20% of the site will be treated with the herbicide.

Local land managers are encouraged to take a triage approach to managing invasive species by prioritizing threats, needs and approach. High priority is given to areas with high ecological values and where control is feasible. Work around these priority areas first addressing small outliers and then moving toward the core of the infestation. The next priority is to address small infestations of high threat species anywhere they can be found and to use the most effective means possible for their control. Once these high priority threats are addressed, land managers

2. Specified controls to mitigate the hazard

should address lower priority areas where control will be effective. The lowest priority are sites with infestations where control is not feasible, at these site land managers should monitor the edge of these sites and implement control efforts to maintain the spread. Throughout this process land managers are encouraged to monitor and learn from the result and to share information gained with others.

3. Program to identify alternatives

The application shall describe the program(s) which are in place in the territory concerned or which will be put in place during the period over which the derogation will be applicable, designed to identify alternative pest control methods which do not use highly hazardous pesticides.

Describe the program(s) that are in place to identify alternatives:

The MDNR continually looks for a variety of methods to control invasive species including biological control, mechanical treatment, cultural practices and herbicides. Over the last ten years the MDNR has worked with Michigan State University, providing financial support and on the ground testing, of several biologic controls. For example, working with MSU, two successful biological controls have been developed for purple loosestrife a non-native invasive wetlands forb. Within the MDNR, research and testing is being conducted to evaluate different methods of controlling invasive species. A recent Wildlife Division project, that has been just concluded, looked at two different herbicides and their effectiveness in controlling autumn olive, a non-native invasive upland shrub. These herbicide practices were evaluated in-conjunction with the use prescribed fire as well as evaluating the impact of prescribed fire only. The use of one of the chemicals and prescribed fire provided the best control.

4. Stakeholder support

All applications for derogations shall include evidence that the application is supported by social, environmental and economic stakeholders in the best interests of promoting FSC's goals in the territory concerned. It is the responsibility of the applicant to present this evidence in support of their application (see summary of procedures in Section 8, below).

The level of stakeholder support required will be evaluated taking account of the geographical scope of the derogation, the justification of need, and other factors include in the application such as the strength of the program to identify alternatives, and the level of controls to mitigate the identified hazards.

A written letter of support by the Board of Directors of the FSC National Initiative for the territory concerned shall normally be considered sufficient evidence of national stakeholder support for the application.

Describe the consultation that has taken place and summarise the results:

A letter addressed to DNR stakeholders, was written by Lynne Boyd, the DNR Division Chief for Forest, Minerals, and Fire Management Division. The letter is dated July 31, 2007, and provides background information in regard to the derogation process and procedure, a list of chemicals for which derogation is being requested, a web address where the actual derogation applications can be viewed, and contact information for the submission of comments. The stakeholder consultation period began August 1, 2007 and ended September 16, 2007. The letter was on DNR letterhead, and its content is shown below:

Dear Stakeholder:

SUBJECT: Opportunity to Comment on Department of Natural Resources Application for Pesticide Derogation (Temporary Exemption) to the Forest Stewardship Council International

The 3.9 million acre Michigan State Forest System is certified by two forest certification systems, the Sustainable Forestry Initiative (SFI) and the Forest Stewardship Council (FSC). The FSC closely scrutinizes the use of pesticide products on certified land, and maintains a list of chemicals that are prohibited from use unless special temporary permission is requested from and granted by the FSC (this is referred to as "derogation") to use them.

The Department of Natural Resources (DNR) is requesting derogation to utilize five pesticides currently on the FSC list. Consistent with FSC policy requirements, we are seeking comments from interested parties on our request for pesticide derogation.

The formal derogation requests, which include detailed information about proposed use of the chemical, are posted on the DNR Forest Certification web site at: http://www.michigan.gov/dnr/0,1607,7-153-30301 33360---,00.html. The following five chemicals are in our derogation request to the FSC:

- Hexazinone (Velpar) one of the most widely used forestry herbicides in the United States. It has been in use for more than 30 years to control brush and weeds during the establishment of new stands of trees.
- 2. 2,4-D, 2-ethylhexyl ester used for habitat restoration and control of invasive exotic plants. It provides a management tool for some difficult to control species.
- 3. 2-(2,4-DP), dma salt (= dichlorprop, dma salt) used for habitat restoration and

- control of invasive exotic plants. It provides a management tool for some difficult to control species.
- 4. Dicamba, dma salt used for habitat restoration and control of invasive exotic plants. It provides a management tool for some difficult to control species.
- 5. Diflybenzuron (Dimlin) used on a limited scale to protect young red pine plantings from Red-headed pine sawfly.

All five of these pesticides have been approved for use by the Environmental Protection Agency, and if used, would be applied according to label guidelines and only by licensed applicators. The use of these chemicals will be minimized, applied on a limited number or acres, and only used when necessary as part of an integrated pest management program to achieve defined management objectives. When use of chemical pesticides is necessary, we select the least toxic, least environmentally persistent, narrowest spectrum products that provide cost effective control and are labeled for the target species.

The DNR invites your comments on use of these pesticides. Please submit your comments by September 16, 2007 to Dennis Nezich, Forest Certification Specialist at Marquette Operations Service Center, 1990 US 41 South, Marquette, MI 49855 or by email at nezichd@michigan.gov. We will compile all comments, include them in the derogation requests, and forward them to the FSC. If you have questions, please feel free to contact Mr. Nezich at 906-228-5245.

Sincerely,

Lynne Boyd, Chief Forest, Mineral, and Fire Management 517-373-1246

Stakeholder consultation actions:

- 1. Lynne Boyd's letter was sent to members of the Michigan Forest Management Advisory Committee on August 12. 2007, following discussion at their August 1, 2007 meeting. The Forest Management Advisory Committee (FMAC) is a 19-member committee whose members are appointed by the Director of the Michigan Department of Natural Resources (DNR). The committee's role is to assist the DNR in balancing the environmental, social and economic issues surrounding forest management. The committee members range from timber producers to university representatives to environmental interests, and assist the DNR with management problems, opportunities and challenges related to Michigan's state forests.
- 2. A prominent notice was posted on the opening page of the Department of Natural Resources internet web site for the entire public input period that began on August 1, 2007 and ended September 16, 2007. This notice, which is displayed below, was linked to Lynne Boyd's stakeholder letter and to copies of the chemical derogation applications.

DNR Asks for Stakeholder Comments on Chemical Derogation
The Department of Natural Resources (DNR) is requesting derogation to
utilize five pesticides. Consistent with the Forest Stewardship Council
(FSC) policy requirements, we are seeking comments from interested
parties on the applications for pesticide derogation.

- 3. Lynne Boyd's letter was sent to members of the Michigan's Sustainable Forestry Initiative, Statewide Implementation Committee. Membership includes representation from forest industry, forestry consultants, universities, forestry associations, and loggers.
- 4. Lynne Boyd's letter was sent to right-of-way maintenance contractors who have held permits for application of chemicals on state forest land.
- 5. A message was posted on the Enviro-Mich Listserv. Enviro-Mich is sponsored by the Sierra Club Mackinac chapter to serve the needs of the citizen environmental and conservation community and citizen's grass roots organizations in Michigan. Enviro-Mich functions like a discussion forum and an automatic e-mail distribution system. Any person who is subscribed to Enviro-Mich can send e-mail to the Enviro-mich address and it will be automatically distributed to the entire list. Cara Boucher, Resource Management Section Leader for the DNR FMFM Division, submitted a notice on September 7, 2007. The notice was titled "Request for public input on Michigan DNR's FSC chemical derogation applications", and included the content of Lynne Boyd's stakeholder letter.
- 6. A request was submitted to the Michigan Environmental Council (MEC) by Cara Boucher, Section Leader in FMFM Division, to forward Lynne Boyd's letter to MEC member organizations for their information. The MEC provides a collective voice for the environment at the local, state, and federal levels. MEC works with 75 member groups and their collective membership. The request was originally made on August 1, 2007. Subsequent contact with MEC on September 11, 2007 revealed that a notice had not been forwarded to members. MEC requested an extension of time for members to comment on the derogation applications. This was granted, with comments due by September 20, 2007.
- 7. An article appeared in the Detroit Free Press on September 6, 2007 titled "Your questions answered: Defining DNR's derogation". This article was written by Free Press Outdoor Writer Eric Sharp (see http://www.freep.com/apps/pbcs.dll/article?AID=2007709060405). The article below includes background information about the FSC derogation process, noted the chemicals for which derogation is sought, and provided DNR contact information for submission of comments. The Free Press also provides an opportunity for reader comments (a single comment was submitted to the Free Press web site).

Your questions answered: Defining DNR's derogation

September 6, 2007

BY ERIC SHARP

FREE PRESS OUTDOORS WRITER

Several readers e-mailed to ask about an item on the Department of Natural Resources Web site that asks for public comment on the "derogation" of five chemicals for use in forestry.

Derogation simply means deviating from a standard and using something in a way that it is not used normally or hasn't been used before.

Dennis Nezich, who works in the Marquette office of the DNR's Forestry, Minerals and Fire Control division, said the chemicals are needed to control the growth of brush, invasive species and a tree-damaging insect.

The reason the state must seek public input is that three years ago the DNR had its forestry program certified by an international body called the Forest Stewardship Council, and the FSC must approve the use of the chemicals.

Part of that approval process requires a chance for stakeholders and members of the public to comment.

Michigan has 3.9 million acres of state forests, and the forestry programs are overseen by both the Sustainable Forestry Initiative and the FSC. The latter organization is especially concerned with things such as the use of chemicals.

"This is an opportunity to take a closer look and ensure that there isn't an undue risk in using the chemicals this way," Nezich said. "The plan will be to use them on a limited basis if it's approved by the FSC."

The DNR didn't join the Forest Stewardship Council just for the cachet. Companies around the world are thinking greener, and many of them won't buy forest products unless they have the FSC seal of approval.

"It's not just in Europe," Nezich said. "A good example is Time Warner. They demand FSC certification," and Time Warner buys a lot of pulpwood to make paper.

Three of the five chemicals will be used to kill invasive species such as knapweed, garlic mustard and autumn olive, and although the chemicals are approved by the U.S. Environmental Protection Agency, the FSC sets a tougher standard.

But it allows the testing to be done by for-profit private agencies, and Marvin Roberson, a forest ecologist for the Sierra Club who lives in Marquette, said that's the suspect link in the chain of protection.

"The Forest Stewardship Council was formed in response to deforestation in the rain forests. It's a good organization. But it hires third-party people to do the certifications. Some of them boast that they've had 100% success in getting (clients certified). That makes you wonder.

"I also am concerned about the standards they use. The standard they're using for Michigan says that the state doesn't have (forest) plantations. But we do -- we have thousands of acres of jack pine plantations."

But overall, Michigan getting FSC certification has been a good thing.

"I'd rather be complaining about a bad statewide standard than having no standard at all," Roberson said.

The chemicals to be derogated are hexazinone, Dimilin, dicamba, 2,4-D dimethylamine salt and 2,4-D 2-ethylexter ester. Members of the public can comment by writing Nezich at the DNR, Marquette Operation Center, 1990 U.S. 41 South, Marquette, MI 49855, or by sending e-mail to him at nezichd@michigan.gov.

Stakeholder Comments received (in chronological order):

1. Izaak Walton League of America, August 13, 2007:

"The Michigan Division of the Izaak Walton League of America supports the Michigan Department of Natural Resources application for pesticide derogations for use of the five pesticides listed in Lynne Boyd's letter of July 31, 2007. Those pesticides include: Hexazinone, 2-3-D, 2-ethylhexyl ester, 2-(2,4_DP) Dicamba, and Diflybenzuron."

"We recognize in some instances there are no good substitutes or the substitutes may be cost prohibitive. As long as the pesticides remain registered for use by the EPA and the Michigan Department of Agriculture, we believe their use should be continued in the manner outlined in Ms. Boyd's letter"

2. Ken Rauscher, Director, Pesticide and Plant Pest Management Division, Michigan Department of Agriculture, August 15, 2007:

"On behalf of the Michigan Department of Agriculture (MDA), Pesticide and Plant Pest Management Division, I am providing the following comments on the Department of Natural Resources Application for Derogation (temporary exemption) to the Forest Stewardship Council International.

MDA is the state agency responsible for pesticide use regulation, including the annual registration of pesticides for use in Michigan. Your request for the use of the 5 pesticides; Hexasinone (Velpar); 2,4-D, 2-ethylhexyl ester; 2-(2,4-DP), dma salt (=dichlorprop, DMA salt); Dicamba, dma salt; and Diflubenzuron (dimilin) under the pesticide derogation application is supported by MDA under the following conditions:

- 1. The specific pesticide chosen is registered for use in Michigan. Currently there are numerous pesticides registered in Michigan that contain the 5 pesticide active ingredients listed in the application.
- 2. Any pesticide use occurs in accordance with label use directions and in compliance with the provisions of the Natural Resources and Environmental Protection Act, Act 451, Part 83, as amended and rules promulgated thereunder.
- Pesticide applicators are certified or registered in accordance with the Natural Resources and Environmental Protection Act, Act 451, Part 83, as amended, and the application of any restricted use pesticide occurs by or under the direct supervision of a Michigan certified applicator.
- 4. Any pesticide application contract issued under the application will only be awarded to a business licensed by MDA for the commercial application of pesticides in all applicable categories of use in accordance with the Natural Resources and Environmental Protection Act, Act 451, Part 83, as amended.

If you have any questions regarding the above comments, please contact Brian Rowe, Pesticide Section Manager, at (517) 373-4905."

3. Jerome Barry, President, Owen Specialty Services, Inc (OSS), August 16, 2007:

"OSS has been engaged in habitat restoration and maintenance since inception in 1993. After review of the intended use for the DNR application for pesticide derogation, I am in favor of said derogation and would probably recommend that the list of products be expanded to include some other products. I believe that there is more than enough Data available from studies and all related industry to show that "Best Management Practices" often must include the qualified and judicious use of these named products to control, and manage specific plant species in the forest population to restore habitat, maintain habitat, limit competition which would harm the plants being established, eliminate invasive species and overall help maintain the healthy diversity of the forest."

4. Mr. Richard Phillips, August 25, 2007:

"The Deregulation of Pesticides is not a good thing. Here in Oakland/Wayne Counties and southeastern Michigan where I grew up in the 60's and 70's I remember all of the bugs that were around, some bit, some ugly and some so beautiful a picture couldn't do them justice. Most all are gone from this region because of the use of pesticides, some bugs were killed

off from the homeowners spraying other from more commercial concentrations sprayed from the air. The eco-ness was effected too, birds, fish, mammals-some couldn't reproduce without deformities! The dragon flies and lighten bugs are only now making a come back to most of this area along with the Hawks. Lets not make the same mistake twice."

5. Mr. Fred Kochis, General Foreman, State Certified Pesticide Applicator, September 7, 2007:

"After reviewing the derogation application information on the MDNR website, Thunder Bay Tree Service, LLC fully supports your request for derogation of the five materials listed:

- 1. Hexazinone (Velpar)
- 2. 2,4-D, 2-ethylhexyl ester
- 3. 2-(2,4-DP), dma salt (= dichlopprop, dma salt)
- 4. Dicamba, dma salt
- 5. Diflybenzuron (Dimlin)

Any integrated pest management program relies on the widest possible assortment of safe and effective methods of pest control. As such, we believe the proper use of these five materials can contribute to a safe and effective I.P.M. program for the establishment and maintenance of Michigan's state forests."

6. Reader comment re: Detroit Free Press article "Your questions answered: Defining DNR's derogation", September 8, 2007:

"If I'm not mistaken, add 2-4-5-T to it and you have agent orange, which by the way was the best broadleaf herbicide I'd ever used. If the Defense Department had used this product as it was directed, the V.A. wouldn't be having the problems it has now. I would think the better the public and private sectors knew about these chemicals, the better off we'd be, and this is an excellent way to do so."

7. Mr. Lee Jackson, Smurfit-Stone Container, Ontonagon, MI, September 11, 2007:

"These selected chemicals are industry standards that should be available for use on state of Michigan lands."

8. Mr. Charles Cubbage, Enviro-mich list serv comment, September 15, 2007:

The following note was posted on Enviro-mich by Mr. Charles Cubbage on September 14, 2007, following some correspondence with the DNR and after additional information was posted on Enviro-mich by the Michigan DNR:

"Thanks for posting the added information. I should imagine that among EMers there are those who would appreciate a note when DNR makes such applications. The transparency is appreciated!"

Contingency plan to eliminate use of the pesticide during the derogation period Derogations shall normally be issued for a five-year period. There is a presumption against renewal at the end of this five-year period unless it can be clearly demonstrated that the program to identify alternatives has been fully implemented but has failed to identify an acceptable alternative in the available time.

Forest managers seeking certification under an approved derogation should therefore ensure that they have a contingency plan in place to eliminate use of the pesticide prior to the end of the derogation period. If a derogation is not renewed, the continued use of a highly hazardous pesticide after the expiry of the derogation would be considered a major non-compliance and would lead to the withdrawal of the certificate.

As a condition of use of a derogated pesticide, forest managers shall record quantitative and qualitative information about their use of such a pesticide, and this information shall be included in the certification body's evaluation reports and in all subsequent surveillance reports.

Compliance with these requirements would need to be demonstrated by an applicant for certification at the Forest Management Unit (FMU) level and be verified by the certification body prior to the issue of a certificate. However, this evaluation is independent of the decision to issue a derogation for use of a pesticide over a geographical area.

References¹

- Environmental Protection Agency. 1998. Tolerances and exemptions from tolerances for pesticide chemicals in food. Part 180 *in* Code of Federal Regulations Title 40, Volume 15 Protection of the Environment.
- Foster, R. K. and R. B. McKercher. 1973. Laboratory incubation studies of chlorophenoxyacetic acids in chernozemic soils. Soil Biol. Biochem. 5:333-337.
- Han, S. O. and P. B. New. 1994. Effect of water availability on degradation of 2,4-dichlorphenoxyacetic acid (2,4-D) by soil microorganisms. Soil Biol. Biochem. 26(12):1689-1697.
- Hayes, H. M., R. E. Tarone, K. P. Cantor, C. R. Jessen, D. M. McCurnin, and R. C. Richardson. 1991. Case-control study of canine malignant lymphoma: positive association with dog owner's use of 2,4-dichlorophenoxyacentic acid herbicides. J. Nation. Cancer Inst. 83:1226-1231.
- Ibrahim, M. A., G. G. Bond, T. A. Burke, P. Cole, F. N. Dost, P. E. Enterline, M. Gough, R. S. Greenberg, W. E. Halperin, E. McConnell, I. C. Munrun, J. A. Swendberg, S. H. Zahm, and J. D. Graham. 1991. Weight of the evidence on the human carcinogencity of 2,4-D. Environ. Health Perspect. 96:213-222.
- Johnson, R. A., and E. V. Wattenberg. 1996. Risk assessment of phenoxy herbicides: an overview of the epidemiology and toxicology data. Chapter 3 in Biological and Economic Assessment of Benefits from Use of Phenoxy Herbicides in the United States. O. C. Burnside ed. U.S.D.A. National Impact Assessment Program. Special NAPIAP Report # 1-PA-96.
- Johnson, W. G., T. L. Lavy, and E. E. Gbur. 1995A. Sorption, mobility, and degradation of triclopyr and 2,4-D and four soils. Weed Sci. 43:678-684.

¹ Much of the reference material used in this paper is from Mandy Tu, Callie Hurd, & John M. Randall, 2001, Weed Control Methods Handbook: Tools and Techniques for Use in Natural Areas. The Nature Conservancy, Global Invasive Species Initiative

- McCall, P. J., S. A. Vrona, and S. S. Kelley. 1981. Fate of uniformly carbon-14 ring labeled 2,4,5-Trichlorophenoxyacetic acid and 2,4-Dichlorophenoxyacetic acid. j. Agric. Food Chem. 29:100-107.
- Mullison, W. R., and G. G. Bond. 1991. Epidemiology and toxicology of 2,4-D. Weed Tech. 5:898-906.
- Newton, M., F. Roberts, A. Allen, B. Kelpsas, D. White, and P. Boyd. 1990. Deposition and dissipation of three herbicides in foliage, litter, and soil of brushfields of Southwest Oregon. J. Agric. Food Chem. 38:574-583.
- Ou, L-T. 1984. 2,4-D degradation and 2,4-D degrading microorganisms in soils. Soil Sci. 137(2):100-107.Parker, L. W. and K. G. Doxtader. 1983. Kinetics of the microbial degradation of 2,4-D in soil: effects of temperature and moisture. J. Environ. Qual. 12(4):553-558.
- Que Hee, S. S., and R. G. Sutherland. 1981. The Phenoxyalkanoic Herbicides, Volume I: Chemistry, Analysis, and Environmental Pollution. CRC Press, Inc., Boca Raton, Florida. 319 pgs.
- Sandmann, E. R. I. C., M. A. Loos, and L. P. van Dyk. 1988. The microbial degradation of 2,4-Dichlorophenoxyacetic acid in soil. Reviews Environ. Contam. Toxicol. 101:1-53.
- Shearer, R., 1980. Public health effects of the aquatic use of herbicides 2,4-D, dichlobenil, endothall and diquat. Section I *in* Literature Reviews of Four Selected Herbicides: 2,4-D, dichlobenil, diquat & endothall. Shearer R., and M. Halter, eds.
- Smith, A. E. 1988. Transformations in soil. Chapter 6 *in* Environmental Chemistry of Herbicides, Volume I. R. Groves ed. CRC Press, Boca Raton, Florida. 207 pgs.
- Thomas, M. L. H., and J. R. Duffy. 1968. Butoxyethanol ester of 2,4-D in the control of eelgrass (*Zostera marina L.*) and its effects on oysters (*Crassostrea viginica Gmelin*) and other benthos. Proc. Northeast. Weed Control. Conf. 22:186
- Tu, M., Hurd, C. & J.M. Randall. 2001. Weed Control Methods Handbook, The Nature Conservancy, http://tncweeds.ucdavis.edu, version: April 2001
 Veeh, R. H., W. P. Inskeep, A. K. Camper. 1996. Soil depth and temperature effects on microbial degradation of 2,4-D. J. Environ. Qual. 25:5-12.
- Wang, Y-S., J-H. Yen, Y-N. Hsieh, and Y-L. Chen. 1994a. Dissipation of 2,4-D, glyphosate, and paraguat in river water. Water Air Soil Pollut. 72:1-7.
- Wang, Y-S., C-G. Jaw, and Y-L. Chen. 1994b. Accumulation of 2,4-D and glyphosate in fish and water hyacinth. Water Air Soil Pollut. 74:397-403.
- Wilson, R. D., J. Geronimo, and J. A. Armbruster. 1997. 2,4-D dissipation in field soils after applications of 2,4-D dimethylamine salt and 2,4-D 2-ethylhexyl ester.
- World Health Organization. 1984. 2,4-Dichlorophenoxyacetic acid (2,4-D), Environmental Health Criteria 29. United Nations Environment Programme, Geneva. 151 pgs.